

Naloxone Distribution and Training



Cuyahoga County Board of Health (CCBH) Overdose Data to Action (OD2A) Initiative

Highlights from OD2A Evaluation findings on naloxone distribution Year One (September, 2019 - August, 2020)

The evaluation for OD2A examines naloxone distribution and training programs, naloxone is a medication administered to rapidly reverse an opioid overdose. It is also known by the brand name, Narcan®.

Each naloxone kit contains two doses (4 mg ea.)



Total Kits Distributed

4,804

Year 1

High Burden Overdose Areas

Zip code data from the Cuyahoga County Medical Examiner's Office, EpiCenter, Ohio EMS, and the Cleveland Division of Police highlight 5 Cuyahoga County zip codes whose residents may benefit from greater access to naloxone.

44109

44102

44111


44105

44103

Overdose Response Training

Overdose response training includes free education on opioid overdose risks and how to recognize a potential overdose and administer naloxone.

MetroHealth provides training to law enforcement, service entities (such as homeless shelters and clinics), and lay responders. Lay responders receive one-on-one training.



Law Enforcement

Trainings: 3

Participants: 48



Service Entities


Trainings: 34

Participants: 202



Lay Responders

Participants: 955




Kit Distribution

Lay responders who received prior overdose response training can also visit distribution partners to receive additional naloxone kits.

MetroHealth Project DAWN Distribution Partners:

- Cuyahoga County Board of Health
- Circle Health Services
- Cleveland Emergency Medical Services
- Cuyahoga County Corrections Center
- Hispanic Urban Minority Alcoholism and Drug Abuse Outreach Program
- Project DAWN Expanded Mobile Unit
- Thomas McCafferty Health Center

Prepared by the Begun Center for Violence Prevention Research and Education, Jack, Joseph and Morton Mandel School of Applied Social Sciences, CaseWestern Reserve University, Evaluation Partner for the OD2A Initiative.



This publication was supported by grant, 5-NU17CE925005-02-00, funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.